Species Comparison Table of Freshwater Mussels of New Jersey

| Freshwater Mussel Species | Lateral Hinge Teeth | Pseudocardinal Teeth | Shape | Maximum Size (inches) | Periostracum Traits | Nacre Color | Other Traits |
|---------------------------|---------------------------------------|------------------------------------|---|-----------------------------|--|--|---|
| Dwarf Wedgemussel | 1 left : 2 right | 2 left : 1 right | elongate triangular | 1.5 | yellowish to blackish | bluish-white | posterior ridge |
| Triangle Floater | absent | 2 left : 1 right robust | triangular inflated | 3 | yellowish green to black; | whitish | green rays in young |
| Brook Floater | absent | 1 left : 1 right small | sub-trapezoidal to reniform - inflated | 3 | yellowish green to greenish brown | bluish white to pale orange | posterior sculpturing orange foot ofter |
| Eastern Elliptio | 2 left : 1 right | 2 left : 1 right well developed | variable - usually sub-trapezoidal | 5 | variable - tan with green rays to black | variable - white pink, gold, purple | green rays in young highly variable |
| Eastern Floater | absent | absent | elongate with rounded ventral margin | 7+ | yellowish green to brown | bluish white | shell thin and fragile banding in youn |
| Alewife Floater | absent | absent | elongate with rounded ventral margin | 7 | yellowish brown variable | white to pale pink | anterior ventral she thicker than res |
| Eastern Pondmussel | 2 left : 1 right thin and delicate | 0 | narrow and elongate compressed | 6 | greenish black to dark brown or black | bluish white | periostracum extend beyond ventral margi |
| Tidewater Mucket | 2 left : 1 right | 2 left : 2 right | ovate inflated | 5 | yellowish to greenish brown | pinkish or salmon | pseudocardinals thi well anterior to bea |
| Yellow Lampmussel | 2 left : 1 right | 2 left : 2-3 right | ovate inflated | 5 | bright yellow to yellowish brown | white | pseudocardinals robus located under bea |
| Eastern Lampmussel | 2 left : 1 right | 2 left : 2-3 right | subovate to ovate slightly inflated | 6 | yellowish green to brownish black | white to pinkish | rays usually prominer |
| Green Floater | 2 left : 1 right vestigal | 2 left : 1 right | ovate to subtrapezoidal variable | 2.5 | yellow to brownish green - thin | bluish white | left valve with interdentation tooth, dark green ray |
| Creeper | absent | swollen knob on each valve | subovate to subtrapezoidal | 4 | yellowish to dark brown or black | white to bluish white | fine rays may be presen |
| Paper Pondshell * | absent | absent | moderately elongate hinge line straight | 4 | greenish to brown - very thin | white to bluish white | beak depressed below hingeline |
| Lilliput * | 2 left : 1 right | 2 left : 1-2 right | elliptical or wedge shaped, inflated | 1.5 | dark green to black | iridescent white | clothlike texture sculpturing when live |
| Chinese Pond Mussel* | absent | absent | ovate-subovate (juveniles) to elongate (adults) can be very inflated when older | 12 | light green (juveniles) to brown (adults)- variable | irridescent white to bluish whte | juveniles with thin, fragile shel and sail-like projection or dorsal margir |

red notes are diagnostic * Introduced species